

Coding Reference Guide Measure Year 2025

Statin Use in Persons with Diabetes (SUPD)



Measure Description

The percentage of Medicare Part D beneficiaries, men and women 40-75 years old, who were dispensed at least two diabetes medication fills on different dates of service and received a statin medication fill between January 1, 2025, and December 31, 2025.

Members included in this measure must have their first fill of diabetes medication at least 90 days before the end of the measurement year or end of the enrollment episode.

Statin Medications and Combinations (Oral Formulations Only):

- Atorvastatin with/without Amlodipine, Ezetimibe
- Fluvastatin
- Lovastatin with/without Niacin
- Pitavastatin
- Pravastatin
- Rosuvastatin with/without Ezetimibe
- Simvastatin with/without Ezetimibe, Niacin

If your patient is unable to use a statin due to one of the following conditions, a claim must be submitted annually using appropriate ICD-10 codes. Please make sure to code using one of the codes below:

The following will exclude Members from this measure:	
Lactation and Fertility	ICD-10: O91.03, O91.13, O91.23, O92.03, O92.13, O92.5, O92.70, O92.79, Z39.1
Pregnancy	ICD-10: O00.101, O00.102, O00.109, O00.111, O00.112, O00.119, O00.201, O00.202, O00.209, O00.211, O00.212, O00.219, O00.80, O00.81, O00.90, O00.91, O09.00, O09.01, O09.02, O09.03, and 1,682 additional codes
End-Stage Renal Disease (ESRD) or Dialysis	ICD-10: I12.0, I13.11, I13.2, N18.5, N18.6, N19, Z91.15, Z99.2
Cirrhosis	ICD-10: K70.30, K70.31, K71.7, K74.3, K74.4, K74.5, K74.60, K74.69
Polycystic Ovary Syndrome (PCOS)	ICD-10: E28.2
Pre-Diabetes	ICD-10: R73.03, R73.09
Rhabdomyolysis and Myopathy Conditions	ICD-10: G72.0, G72.89, G72.9, M60.80, M60.819, M60.829, M60.839, M60.849 M60.859, M60.869, M60.879, M60.9, M62.82
Hospice Enrollment	

Tips and Best Practices to Help Improve Performance

- Include the following topics in your discussions with patients who are prescribed statin medications:
 - Importance of statin medication adherence and lifestyle modifications (diet, exercise, weight loss)
 - Side effects such as muscle pain and what to do if they occur
 - What to do about missed doses
 - Identify barriers that would impact adherence to statin therapy
- Once the patient demonstrates that they tolerate statin therapy, prescribe a 90-day supply
- Consider using alerts and flags within the Electronic Health Record (EHR) for screening tests
- Discuss the importance of early detection and encourage preventive care
- Document preventive care, medical, and surgical history in the medical record with specific date and results
- Use correct diagnosis and procedure codes and submit claims and encounter data in a timely manner
- Stock patient areas and exam rooms with educational materials such as those found here:
 - [Diabetes | CDC](#)
 - [Medical Health - MVP Health Care](#)